

AD-A136 595

THE TOTAL FIELD IN COLLECTIVE BREMSSTRAHLUNG IN A  
NONEQUILIBRIUM RELATIVISTIC BEAM-PLASMA SYSTEM (U) HARRY  
DIAMOND LABS ADELPHI MD H E BRANDT SEP 83 HDL-TR-1996


1/1

UNCLASSIFIED

F/G 28/9.

NL

END



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER HDL-TR-1996	2. GOVT ACCESSION NO. AD-A136545	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Total Field in Collective Bremsstrahlung in a Nonequilibrium Relativistic Beam-Plasma System		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Howard E. Brandt		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Harry Diamond Laboratories 2800 Powder Mill Road Adelphi, MD 20783		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Program Ele: 61101A
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Materiel Development and Readiness Command Alexandria, Va 22333		12. REPORT DATE September 1983
		13. NUMBER OF PAGES 34
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES HDL Project: A10225 DRCMS Code: 61110191A0011 DA Project: 1L161101A91A		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Radiation                      Relativistic electron beams Microwaves                    Radiative instability Plasma turbulence            Bremsstrahlung Relativistic plasma          Beam-plasma systems Plasma physics               Nonlinear plasmas		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An expression is obtained for the total electric field associated with collective bremsstrahlung in a nonequilibrium relativistic beam-plasma system in the case of a slowly varying, nearly spatially independent background distribution with no external fields. The field is assumed to consist of an unperturbed regular part corresponding to the nonradiative part of the field of the particles and a stochastic part corresponding to the bremsstrahlung radiation field. The relations derived here are needed in calculations of collective bremsstrahlung processes and the conditions for the occurrence of bremsstrahlung radiative instability in relativistic beam-plasma systems.		

DD FORM 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

# CONTENTS

	<u>Page</u>
1. INTRODUCTION .....	5
2. DISPERSION RELATION FOR THE BREMSSTRAHLUNG FIELD .....	6
3. THE STOCHASTIC PROPERTIES OF THE BREMSSTRAHLUNG FIELD .....	11
4. THE FIELD DUE TO THE PERTURBED TEST PARTICLE TRAJECTORY .....	13
5. THE FIELD DUE TO THE DYNAMIC POLARIZATION CURRENT .....	18
6. THE TOTAL FIELD .....	22
7. CONCLUSION .....	23
LITERATURE CITED .....	24
DISTRIBUTION .....	27

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## 1. INTRODUCTION

In a nonequilibrium relativistic beam-plasma system with a slowly varying, nearly spatially independent background distribution without external fields, the total field, excited in the beam-plasma and associated with a collective bremsstrahlung process involving a test particle, has two parts. First, it has a regular part with a relatively slowly varying phase associated with the nonradiative field of the test particle and its induced dynamic polarization, and it also has a stochastic part with an irregular rapidly varying phase associated with the bremsstrahlung radiation field.<sup>1</sup> Thus, the Fourier transform of the total field involved in the bremsstrahlung process is given by

$$\vec{E}_k = \vec{E}_k^R + \vec{E}_k^{st}, \quad (1)$$

where  $\vec{E}_k^R$  and  $\vec{E}_k^{st}$  are the regular nonradiative component and the stochastic bremsstrahlung component, respectively. The regular part,  $\vec{E}_k^R$ , of this field is assumed to consist of four parts in sufficient approximation; namely,

$$\vec{E}_k^R = \vec{E}_k^{(1)} + \vec{E}_k^{(2)} + \vec{E}_{dpk}^{(1)} + \vec{E}_{dpk}^{(2)}. \quad (2)$$

Here  $\vec{E}_k^{(1)}$  is the self-field associated with the unperturbed motion of the relativistic bare test particle, ignoring its induced dynamic polarization. The field  $\vec{E}_k^{(2)}$  is the field associated with the perturbation in the bare particle motion. The fields  $\vec{E}_{dpk}^{(1)}$  and  $\vec{E}_{dpk}^{(2)}$  are fields produced by the dynamic polarization current induced by the test particle to second and third order in the total field, respectively.

In sections 2 through 5, the dispersive characteristics and stochastic properties of the bremsstrahlung field are reviewed, the field due to the perturbed bare test particle trajectory is calculated from the associated current, and the dynamic polarization current is used to calculate the field which it induces. In section 6 an expression for the total field is obtained, and in section 7 the results are summarized. Expressions for these fields are needed in calculations of the total nonlinear force on a relativistic test particle involved in collective bremsstrahlung in a nonequilibrium beam-plasma

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

system.<sup>1,\*†</sup> The latter is needed in calculations of collective bremsstrahlung processes and the conditions for the occurrence of a bremsstrahlung radiative instability.<sup>1-4,\*†‡</sup>

## 2. DISPERSION RELATION FOR THE BREMSSTRAHLUNG FIELD

In zeroth approximation the dispersive properties of the bremsstrahlung radiation field  $\vec{E}_k^{st}$  are governed by the poles of the linear photon Green's function  $G_{mn}(k)$  for the beam-plasma. The latter determines the mode dispersion relations. Thus,

$$\vec{E}_k^{st} = \vec{E}_k^{\sigma(0)}, \quad (3)$$

where the electromagnetic waves  $\vec{E}_k^{\sigma(0)}$  of mode  $\sigma$  satisfy the following dispersion relation:

$$|G_{mn}^{-1}(k)| = 0. \quad (4)$$

The linear photon Green's function  $G_{mn}(k)$  follows from the linear Maxwell's equations in the beam-plasma system, namely,

$$\vec{\nabla} \times \vec{H} = \vec{j} + \frac{\partial \vec{D}}{\partial t}, \quad (5)$$

and

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

<sup>2</sup>A. V. Akopyan and V. N. Tsytovich, *Transition Bremsstrahlung of Relativistic Particles*, *Zh. Eksp. Teor. Fiz.*, 71 (1976), 166 [*Sov. Phys. JETP*, 44 (1976), 87].

<sup>3</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung Instability of Relativistic Electrons in a Plasma*, *Astrofizika*, 13 (1977), 717 [*Astrophysics*, 13 (1977), 423].

<sup>4</sup>V. N. Tsytovich, *Collective Effects in Bremsstrahlung of Fast Particles in Plasmas*, *Comments, Plasma Phys. Conf. Fusion*, 4 (1978), 73.

\*H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-7 (May 1982) (to be published as HDL-TR-1995, 1983).

†H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

‡H. E. Brandt, *Theoretical Methods in the Calculation of the Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-83-6 (April 1983) (to be published as HDL-TR-2009, 1983).

$$\vec{\nabla} \times \vec{E} = - \frac{\partial \vec{B}}{\partial t} . \quad (6)$$

The Fourier decomposition of the fields is given by

$$\vec{E} = \int d\mathbf{k} \vec{E}_k e^{i(\vec{k} \cdot \vec{r} - \omega t)} \quad (7)$$

and

$$\vec{B} = \int d\mathbf{k} \vec{B}_k e^{i(\vec{k} \cdot \vec{r} - \omega t)} , \quad (8)$$

where

$$d\mathbf{k} = d^3\mathbf{k} d\omega . \quad (9)$$

The assumed forms of the constitutive relations for the beam-plasma system are

$$\vec{H}_k = \frac{\vec{B}_k}{\mu_0} , \quad (10)$$

where  $\mu_0$  is the permeability of free space, and

$$D_{ki} = \epsilon_{ij}(k) E_{kj} , \quad (11)$$

where  $\epsilon_{ij}(k)$  is the dielectric tensor for the beam-plasma system, and summation over repeated indices is understood. From equations (5), (10), and (11) it follows that

$$\frac{i}{\mu_0} (\vec{k} \times \vec{B}_k)_i + i\omega \epsilon_{ij}(k) E_{kj} = j_{ki} , \quad (12)$$

where  $\vec{j}_k$  is the Fourier transform of the nonlinear and external current density, and from equation (6)

$$\vec{B}_k = \frac{\vec{k}}{\omega + i\delta} \times \vec{E}_k . \quad (13)$$

Here  $\delta$  is a small imaginary part of the frequency. Next, substituting equation (13) in equation (12), then

$$\frac{i}{\mu_0(\omega + i\delta)} [\vec{k} \times (\vec{k} \times \vec{E}_k)]_i + i\omega \epsilon_{ij} E_{kj} = j_{ki} . \quad (14)$$

But one has the following vector identity,

$$\vec{k} \times (\vec{k} \times \vec{E}_k) = (\vec{k} \cdot \vec{E}_k) \vec{k} - k^2 \vec{E}_k . \quad (15)$$

Substituting equation (15) in equation (14), then

$$\frac{i}{\mu_0(\omega + i\delta)} (\vec{k} \cdot \vec{E}_k k_i - k^2 E_{ki}) + i\omega \epsilon_{ij} E_{kj} = j_{ki} \quad (16)$$

Equation (16) may be rewritten as

$$G_{ij}^{-1} E_{kj} = - \frac{i}{\omega + i\delta} j_{ki} \quad (17)$$

where

$$G_{ij}^{-1} = \frac{1}{\mu_0(\omega + i\delta)^2} (k_i k_j - k^2 \delta_{ij}) + \epsilon_{ij} \quad (18)$$

where  $\delta_{ij}$  is the Kronecker delta. For a spatially isotropic system, the dielectric permittivity tensor is given by<sup>5,6,7</sup>

$$\epsilon_{ij}(\vec{k}, \omega) = \left( \delta_{ij} - \frac{k_i k_j}{k^2} \right) \epsilon_t(\vec{k}, \omega) + \frac{k_i k_j}{k^2} \epsilon_l(\vec{k}, \omega) \quad (19)$$

where  $\epsilon_t$  and  $\epsilon_l$  are the transverse and longitudinal permittivity, respectively. Substituting equation (19) in equation (18) and combining terms, then

$$G_{ij}^{-1} = \left( \epsilon_t - \frac{k^2}{\mu_0(\omega + i\delta)^2} \right) \delta_{ij} - \left( \epsilon_t - \epsilon_l - \frac{k^2}{\mu_0(\omega + i\delta)^2} \right) \frac{k_i k_j}{k^2} \quad (20)$$

Treating equation (17) as a matrix equation, then

$$E_{ki} = - \frac{i}{\omega + i\delta} G_{ij} j_{kj} \quad (21)$$

The matrix  $G_{ij}$  inverse to  $G_{ij}^{-1}$  is easily obtained from equation (20); namely,

$$G_{ij} = \frac{k_i k_j}{k^2 \epsilon_l} + \frac{\omega^2}{\epsilon_t(\omega + i\delta)^2 - \frac{k^2}{\mu_0}} \left( \delta_{ij} - \frac{k_i k_j}{k^2} \right) \quad (22)$$

Checking that equation (22) is in fact the inverse, one notes, using equations (20) and (22), that

<sup>5</sup>A. I. Akhiezer, I. A. Akhiezer, R. V. Polovin, A. G. Sitenko, and K. N. Stepanov, *Plasma Electrodynamics, 1, Linear Theory*, Pergamon Press (1975), 206.

<sup>6</sup>V. N. Tsytovich, *Nonlinear Effects in Plasma*, Plenum Press, New York (1970), 31.

<sup>7</sup>V. N. Tsytovich, *Theory of Turbulent Plasma (Consultants Bureau)*, Plenum Press, New York (1977), 63-65.

$$G_{mi}^{-1} G_{ij} = \left[ \left( \epsilon_t - \frac{k^2}{\mu_0(\omega + i\delta)^2} \right) \delta_{mi} - \left( \epsilon_t - \epsilon_l - \frac{k^2}{\mu_0(\omega + i\delta)^2} \right) \frac{k_m k_i}{k^2} \right] \quad (23)$$

$$\times \left[ \frac{k_i k_j}{k^2 \epsilon_l} + \frac{\omega^2}{\epsilon_t(\omega + i\delta)^2 - \frac{k^2}{\mu_0}} \left( \delta_{ij} - \frac{k_i k_j}{k^2} \right) \right] = \delta_{mj}$$

as it must. Equation (21) can be equivalently written in the following form:

$$E_{kn} = - \frac{i}{\omega + i\delta} G_{mn} j_{km} \quad , \quad (24)$$

since the linear photon Green's function equation (22), for the isotropic plasma is a symmetric tensor. According to equation (4), the poles of this function determine the zeroth order dispersion relation for the bremsstrahlung in the isotropic case.

More generally the dielectric tensor is related to the linear conductivity tensor by

$$\epsilon_{ij} = \epsilon_0 \delta_{ij} + \frac{i \sigma_{ij}(k)}{\omega + i\delta} \quad . \quad (25)$$

Equation (25) follows through a comparison of equation (12) with the vacuum form of Maxwell's equations; namely,

$$\frac{i}{\mu_0} (\vec{k} \times \vec{B}_k)_i + i\omega \epsilon_0 \delta_{ij} E_{kj} = \sigma_{ij}(k) E_{kj} + j_{ki} \quad , \quad (26)$$

where the explicit linear current in terms of the linear conductivity has been introduced on the right,  $j_{ki}$  designates the nonlinear and external contributions to the current, and the vacuum dielectric tensor has been introduced. Thus, moving the first term on the right of equation (26) to the left and comparing with equation (12), then equation (25) follows.

By equation (47) of HDL-TR-1994,\* the linear conductivity tensor  $\sigma_{ij}$  in the beam plasma system is given by

$$\sigma_{ij} = \sum_s e_s^2 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_{si} \left[ \delta_{jm} \left( 1 - \frac{\vec{k} \cdot \vec{v}_s}{\omega + i\delta} \right) + \frac{k_m v_{sj}}{\omega + i\delta} \right]}{i(\omega - \vec{k} \cdot \vec{v}_s + i\delta)} \frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} \quad . \quad (27)$$

---

\*H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

Here,  $s$  is a species label. Substituting equation (27) in equation (25), then the linear dielectric tensor for the beam-plasma system is given by

$$\epsilon_{ij} = \epsilon_0 \delta_{ij} + \frac{1}{\omega + i\delta} \sum_s e_s^2 \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{v_{si} \left[ \delta_{jm} \left( 1 - \frac{\vec{k} \cdot \vec{v}_s}{\omega + i\delta} \right) + \frac{k_m v_{sj}}{\omega + i\delta} \right]}{(\omega - \vec{k} \cdot \vec{v}_s + i\delta)} \frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} \quad (28)$$

in which case equation (18) becomes

$$G_{ij}^{-1} = \frac{1}{\mu_0 (\omega + i\delta)^2} (k_i k_j - k^2 \delta_{ij}) + \epsilon_0 \delta_{ij} + \frac{1}{\omega + i\delta} \sum_s e_s^2 \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{v_{si} \left[ \delta_{jm} \left( 1 - \frac{\vec{k} \cdot \vec{v}_s}{\omega + i\delta} \right) + \frac{k_m v_{sj}}{\omega + i\delta} \right]}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} \quad (29)$$

It should be noted that equations (27), (28), and (29) are in fact symmetric in  $i$  and  $j$  which follows after a simple integration by parts. The zeros of the determinant of the matrix equation (29) must give the dispersive properties of the bremsstrahlung field.

From equation (19) one has that

$$\epsilon_l = \frac{k_i k_j}{k^2} \epsilon_{ij} \quad (30)$$

and

$$\epsilon_t = \frac{1}{2} (\epsilon_{ii} - \epsilon_l) \quad (31)$$

Substituting equation (28) in equations (30) and (31), then in the spatially isotropic case, the longitudinal and transverse dielectric permittivity are given by

$$\epsilon_l = \epsilon_0 + \frac{1}{\omega + i\delta} \sum_s \frac{e_s^2}{k^2} \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{\vec{k} \cdot \vec{v}_s}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \vec{k} \cdot \vec{v}_{ps} f_{ps}^{R(0)} \quad (32)$$

and

$$\epsilon_t = \epsilon_0 + \frac{1}{2(\omega + i\delta)^2} \sum_s \frac{e_s^2}{k^2} \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \left[ k^2 v_{sm} + \frac{k^2 v_s^2 - \omega \vec{k} \cdot \vec{v}_s}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} k_m \right] \frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} \quad (33)$$

If the background distribution functions  $f_{ps}^{R(0)}$  are also isotropic in momentum space, then they depend only on particle energy; namely,

$$f_{ps}^{R(0)} = f_{ps}^{R(0)}(\epsilon_s) \quad (34)$$

where

$$\epsilon_s = (p_s^2 c^2 + m_s^2 c^4)^{1/2} . \quad (35)$$

Then

$$\frac{\partial f_{ps}^{R(0)}}{\partial p_{sm}} = \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} \frac{\partial \epsilon_s}{\partial p_{sm}} = \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} \frac{p_{sm} c^2}{\epsilon_s} = v_{sm} \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} . \quad (36)$$

Substituting equation (36) in equation (28) and simplifying, then for the case of isotropy in momentum space, one has

$$\epsilon_{ij} = \epsilon_0 \delta_{ij} + \frac{1}{\omega + i\delta} \sum_s e_s^2 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_{si} v_{sj}}{(\omega - \vec{k} \cdot \vec{v}_s + i\delta)} \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} . \quad (37)$$

Substituting equation (37) in equations (30) and (31), then in the case of isotropy in momentum space, the longitudinal and transverse dielectric permittivity are given by

$$\epsilon_l = \epsilon_0 + \frac{1}{\omega + i\delta} \sum_s \frac{e_s^2}{k^2} \int \frac{d^3 p_s}{(2\pi)^3} \frac{(\vec{k} \cdot \vec{v}_s)^2}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} \quad (38)$$

and

$$\epsilon_t = \epsilon_0 + \frac{1}{2} \frac{1}{\omega + i\delta} \sum_s e_s^2 \int \frac{d^3 p_s}{(2\pi)^3} \frac{v_s^2 - (\vec{k} \cdot \vec{v}_s)^2 / k^2}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \frac{\partial f_{ps}^{R(0)}}{\partial \epsilon_s} , \quad (39)$$

and the zeros of the determinant of equation (20) along with equations (38) and (39) then give the dispersion relation for the bremsstrahlung in this case.

### 3. THE STOCHASTIC PROPERTIES OF THE BREMSSTRAHLUNG FIELD

As stated earlier, it is to be assumed that the bremsstrahlung field resulting from the particle scattering in the system is stochastic, having an irregular rapidly varying phase.<sup>1</sup> To the needed order, the phase of the stochastic bremsstrahlung field is assumed to change randomly during a typical interaction period, and the time average is assumed to be the same as the average over the stochastic ensemble. In particular it is assumed that for this stochastic field the following averages over the statistical phase distribution apply approximately; namely,

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

$$\langle E_{ki}^{st} \rangle = 0 , \quad (40)$$

$$\langle E_{ki}^{st} E_{k_1j}^{st} \rangle = |E_k^{st}|^2 \delta(k + k_1) e_{ki} e_{k_1j}^* , \quad (41)$$

where  $\vec{e}_k$  is the unit polarization vector for the stochastic field.

Equations (40) and (41) are deduced as follows. Representing the stochastic electric field in terms of its modulus,  $A_k$ , and phase,  $\phi_k$ , one has

$$E_{ki}^{st} = e_{ki} A_k e^{i\phi_k} = e_{ki} A_k (\cos \phi_k + i \sin \phi_k) . \quad (42)$$

Therefore, the average over the statistical phase distribution  $\langle E_{ki}^{st} \rangle$  is given by

$$\langle E_{ki}^{st} \rangle = e_{ki} A_k (\langle \cos \phi_k + i \sin \phi_k \rangle) . \quad (43)$$

But assuming random phase to the needed order, then clearly

$$\langle \cos \phi_k \rangle = \langle \sin \phi_k \rangle = 0 , \quad (44)$$

and equation (43) becomes equation (40). Similarly,

$$\langle E_{ki}^{st} E_{k_1j}^{st*} \rangle = e_{ki} e_{k_1j}^* A_k A_{k_1} \langle e^{i\phi_k} e^{-i\phi_{k_1}} \rangle . \quad (45)$$

But

$$\langle e^{i\phi_k} e^{-i\phi_{k_1}} \rangle = \langle e^{i(\phi_k - \phi_{k_1})} \rangle = \delta_{k,k_1} , \quad (46)$$

since it is vanishing unless  $k = k_1$  and unity when  $k = k_1$ . Substituting equation (46) in equation (45), using the property of the Kronecker delta, denoting  $k_1$  by  $-k_1$ , and using the reality property of the field ( $\vec{E}_{-k}^* = \vec{E}_k$ ), then

$$\langle E_{ki}^{st} E_{k_1j}^{st} \rangle = e_{ki} e_{k_1j}^* A_k^2 \delta_{-k_1,k} . \quad (47)$$

Next, one may define the quantity  $|E_k^{st}|$  in terms of the modulus  $A_k$  by

$$|E_k^{st}|^2 \delta(k + k_1) = A_k^2 \delta_{-k_1,k} . \quad (48)$$

Then substituting equation (48) in equation (47), equation (41) follows.

Furthermore, using equation (3) in equations (40) and (41) one has for the stochastic properties of the bremsstrahlung radiation field the following:

$$\langle E_{ki}^{\sigma(0)} \rangle = 0 , \quad (49)$$

$$\langle E_{ki}^{\sigma(0)} E_{k_1j}^{\sigma(0)*} \rangle = e_{ki} e_{k_1j}^* |E_k^{\sigma(0)}|^2 \delta(k + k_1) , \quad (50)$$

where  $\vec{e}_k^\sigma$  is the unit polarization vector for mode  $\sigma$ .

#### 4. THE FIELD DUE TO THE PERTURBED TEST PARTICLE TRAJECTORY

The trajectory of a relativistic test particle of species  $\alpha$  moving through the beam-plasma system is given by

$$\vec{r}_\alpha = \vec{v}_\alpha t + \Delta \vec{r}_\alpha, \quad (51)$$

where  $\vec{v}_\alpha$  is the unperturbed velocity and  $\Delta \vec{r}_\alpha$  and  $\Delta \vec{v}_\alpha$  describe the perturbation in the trajectory and the velocity, respectively. Provided the conditions for the plasma Born approximation are satisfied--namely, that the relativistic particle momentum be much greater than the electromagnetic impulse received by the particle in a time interval given by the inverse electron plasma frequency--then the perturbed motion is given by equation (36) of HDL-TR-1995:\*

$$\begin{aligned} \Delta \vec{r}_\alpha = & -\frac{e_\alpha}{\gamma_\alpha m_\alpha} \int \frac{d^3 \vec{k} d\omega}{(\omega - \vec{k} \cdot \vec{v}_\alpha + i\delta)^2} \left[ \vec{E}_k \left( 1 - \frac{\vec{k} \cdot \vec{v}_\alpha}{\omega + i\delta} \right) + \vec{k} \frac{\vec{v}_\alpha \cdot \vec{E}_k}{\omega + i\delta} \right] e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t} \\ & + \frac{e_\alpha \vec{v}_\alpha}{\gamma_\alpha m_\alpha c^2} \int \frac{d^3 \vec{k} d\omega}{(\omega - \vec{k} \cdot \vec{v}_\alpha + i\delta)^2} \vec{v}_\alpha \cdot \vec{E}_k e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t}. \end{aligned} \quad (52)$$

Here  $e_\alpha$ ,  $m_\alpha$ , and  $\vec{v}_\alpha$  are the charge, mass, and unperturbed velocity of the test particle, respectively. (For notational convenience, the naught in HDL-TR-1995 is dropped here.) The unperturbed relativistic factor  $\gamma_\alpha$  is given by

$$\gamma_\alpha = \left[ 1 - \left( \frac{v_\alpha}{c} \right)^2 \right]^{-1/2}. \quad (53)$$

The perturbation in the velocity is given by equation (35) of HDL-TR-1995:\*

$$\begin{aligned} \Delta \vec{v}_\alpha = & \frac{ie_\alpha}{\gamma_\alpha m_\alpha} \int d^3 \vec{k} d\omega \left[ \vec{E}_k \left( 1 - \frac{\vec{k} \cdot \vec{v}_\alpha}{\omega + i\delta} \right) + \vec{k} \left( \frac{\vec{v}_\alpha \cdot \vec{E}_k}{\omega + i\delta} \right) \right] \frac{e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t}}{\omega - \vec{k} \cdot \vec{v}_\alpha + i\delta} \\ & - \frac{ie_\alpha \vec{v}_\alpha}{\gamma_\alpha m_\alpha c^2} \int d^3 \vec{k} d\omega \frac{\vec{v}_\alpha \cdot \vec{E}_k e^{-i(\omega - \vec{k} \cdot \vec{v}_\alpha)t}}{\omega - \vec{k} \cdot \vec{v}_\alpha + i\delta}. \end{aligned} \quad (54)$$

The  $i^{\text{th}}$  component of  $\Delta \vec{v}_\alpha$ , equation (54), can be equivalently rewritten as follows:

---

\*H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-7 (May 1982) (to be published as HDL-TR-1995, 1983).

$$\Delta v_{\alpha i} = \frac{-ie_{\alpha}}{\gamma_{\alpha} m_{\alpha}} \int d^3 \vec{k} d\omega \left( \frac{v_{\alpha i} v_{\alpha j}}{c^2} - \delta_{ij} \right) \left[ E_{kj} \left( 1 - \frac{\vec{k} \cdot \vec{v}_{\alpha}}{\omega + i\delta} \right) + k_j \frac{\vec{v}_{\alpha} \cdot \vec{E}_{\vec{k}}}{\omega + i\delta} \right] \frac{e^{-i(\omega - \vec{k} \cdot \vec{v}_{\alpha})t}}{\omega - \vec{k} \cdot \vec{v}_{\alpha} + i\delta} . \quad (55)$$

The current density due to the actual charge of the test particle is then given by

$$\vec{j}_{\alpha} = e_{\alpha} (\vec{v}_{\alpha} + \Delta \vec{v}_{\alpha}) \delta^3(\vec{r} - \vec{v}_{\alpha} t - \Delta \vec{r}_{\alpha}) . \quad (56)$$

Expanding the delta function in equation (56) for small  $\Delta \vec{r}_{\alpha}$ , then to lowest order in  $\Delta \vec{r}_{\alpha}$ ,

$$\vec{j}_{\alpha} = e_{\alpha} (\vec{v}_{\alpha} + \Delta \vec{v}_{\alpha}) [\delta^3(\vec{r} - \vec{v}_{\alpha} t) - \Delta \vec{r}_{\alpha} \cdot \vec{\nabla} \delta^3(\vec{r} - \vec{v}_{\alpha} t)] . \quad (57)$$

Thus, to lowest order in  $\Delta \vec{r}_{\alpha}$  and  $\Delta \vec{v}_{\alpha}$ , one has

$$\vec{j}_{\alpha} = \vec{j}_{\alpha}^{(1)} + \vec{j}_{\alpha}^{(2)} , \quad (58)$$

where

$$\vec{j}_{\alpha}^{(1)} = e_{\alpha} \vec{v}_{\alpha} \delta^3(\vec{r} - \vec{v}_{\alpha} t) \quad (59)$$

and

$$\vec{j}_{\alpha}^{(2)} = e_{\alpha} \Delta \vec{v}_{\alpha} \delta^3(\vec{r} - \vec{v}_{\alpha} t) - e_{\alpha} \vec{v}_{\alpha} \Delta \vec{r}_{\alpha} \cdot \vec{\nabla} \delta^3(\vec{r} - \vec{v}_{\alpha} t) . \quad (60)$$

The Fourier components of these currents at the position of the test particle are given by

$$j_{\alpha ki}^{(1)} = \int \frac{d^3 \vec{r} d\vec{t}}{(2\pi)^4} j_{\alpha i}^{(1)}(\vec{r}, t) e^{-i(\vec{k} \cdot \vec{r} - \omega t)} \quad (61)$$

$$j_{\alpha ki}^{(2)} = \int \frac{d^3 \vec{r} d\vec{t}}{(2\pi)^4} j_{\alpha i}^{(2)}(\vec{r}, t) e^{-i(\vec{k} \cdot \vec{r} - \omega t)} . \quad (62)$$

Substituting equation (59) in equation (61) and integrating over the delta function, then

$$j_{\alpha ki}^{(1)} = \frac{e_{\alpha} v_{\alpha i}}{(2\pi)^3} \int \frac{d\vec{t}}{2\pi} e^{-i(\vec{k} \cdot \vec{v}_{\alpha} - \omega)t} . \quad (63)$$

Next, recognizing the integral form of the delta function appearing in equation (63), then

$$j_{\alpha ki}^{(1)} = \frac{e_{\alpha} v_{\alpha i}}{(2\pi)^3} \delta(\omega - \vec{k} \cdot \vec{v}_{\alpha}) . \quad (64)$$

Next, substituting equation (60) in equation (62) one has

$$j_{aki}^{(2)} = \int \frac{d^3 \vec{r}}{(2\pi)^4} e_{\alpha} \Delta v_{\alpha i} \delta^3(\vec{r} - \vec{v}_{\alpha} t) e^{-i(\vec{k} \cdot \vec{r} - \omega t)} - \int \frac{d^3 \vec{r}}{(2\pi)^4} e_{\alpha} v_{\alpha i} \Delta \vec{r}_{\alpha} \cdot [\vec{v} \delta^3(\vec{r} - \vec{v}_{\alpha} t)] e^{-i(\vec{k} \cdot \vec{r} - \omega t)} . \quad (65)$$

The second integral in equation (65) can be simplified by integration over space by parts and dropping surface terms. Equation (65) becomes

$$j_{aki}^{(2)} = \int \frac{d^3 \vec{r}}{(2\pi)^4} e_{\alpha} \Delta v_{\alpha i} \delta^3(\vec{r} - \vec{v}_{\alpha} t) e^{-i(\vec{k} \cdot \vec{r} - \omega t)} - \int \frac{d^3 \vec{r}}{(2\pi)^4} e_{\alpha} v_{\alpha i} \delta^3(\vec{r} - \vec{v}_{\alpha} t) i \vec{k} \cdot \Delta \vec{r}_{\alpha} e^{-i(\vec{k} \cdot \vec{r} - \omega t)} . \quad (66)$$

Integrating over the delta functions, then

$$j_{aki}^{(2)} = \int \frac{dt}{(2\pi)^4} e_{\alpha} \Delta v_{\alpha i} e^{-i(\vec{k} \cdot \vec{v}_{\alpha} - \omega)t} - i \int \frac{dt}{(2\pi)^4} e_{\alpha} v_{\alpha i} e^{-i(\vec{k} \cdot \vec{v}_{\alpha} - \omega)t} \vec{k} \cdot \Delta \vec{r}_{\alpha} . \quad (67)$$

Next, substituting equations (52) and (55) in equation (67) and integrating over the time, then

$$j_{aki}^{(2)} = \frac{-ie_{\alpha}^2}{\gamma_{\alpha} m_{\alpha}} \left( \frac{v_{\alpha i} v_{\alpha j}}{c^2} - \delta_{ij} \right) \int \frac{d^3 \vec{k}_1 d\omega_1}{(2\pi)^3} \left[ E_{k_1 j} \left( 1 - \frac{\vec{k}_1 \cdot \vec{v}_{\alpha}}{\omega_1 + i\delta} \right) + k_{1j} \frac{\vec{v}_{\alpha} \cdot \vec{E}_{k_1}}{\omega_1 + i\delta} \right] \frac{\delta(\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_{\alpha})}{\omega_1 - \vec{k}_1 \cdot \vec{v}_{\alpha} + i\delta} + \frac{ie_{\alpha}^2}{\gamma_{\alpha} m_{\alpha}} v_{\alpha i} \int \frac{d^3 \vec{k}_1 d\omega_1}{(2\pi)^3} \frac{\left[ \vec{k} \cdot \vec{E}_{k_1} \left( 1 - \frac{\vec{k}_1 \cdot \vec{v}_{\alpha}}{\omega_1 + i\delta} \right) + \frac{\vec{k} \cdot \vec{k}_1 \vec{v}_{\alpha} \cdot \vec{E}_{k_1}}{\omega_1 + i\delta} \right]}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_{\alpha} + i\delta)^2} \times \delta(\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_{\alpha}) - \frac{ie_{\alpha}^2}{\gamma_{\alpha} m_{\alpha} c^2} v_{\alpha i} \int \frac{d^3 \vec{k}_1 d\omega_1}{(2\pi)^3} \frac{\vec{k} \cdot \vec{v}_{\alpha} \vec{v}_{\alpha} \cdot \vec{E}_{k_1}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_{\alpha} + i\delta)^2} \delta(\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_{\alpha}) . \quad (68)$$

Simplifying equation (68), then

$$\begin{aligned}
j_{aki}^{(2)} = & \frac{ie_a^2}{\gamma_a^m} \int \frac{d^3k_1 d\omega_1}{(2\pi)^3} \frac{E_{k_1j}}{\omega_1 + i\delta} \delta(\omega - \omega_1 - \vec{k} \cdot \vec{v}_a + \vec{k}_1 \cdot \vec{v}_a) \\
& \times \left\{ \delta_{ij} - \frac{v_{ai}v_{aj}}{c^2} - \frac{\vec{k}_1 \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta)c^2} + \frac{k_{1i}v_{aj}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta} \right. \\
& \left. + \frac{k_j v_{ai}}{\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta} + \frac{v_{ai}v_{aj} \vec{k} \cdot \vec{k}_1}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta)^2} - \frac{\omega_1 \vec{k} \cdot \vec{v}_a v_{ai}v_{aj}}{c^2(\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta)^2} \right\}.
\end{aligned} \quad (69)$$

Using the property of the delta function to replace  $(\omega_1 - \vec{k}_1 \cdot \vec{v}_a)$  by  $(\omega - \vec{k} \cdot \vec{v}_a)$  in the fourth, fifth, and sixth terms of equation (69), then

$$\begin{aligned}
j_{aki}^{(2)} = & \frac{ie_a^2}{\gamma_a^m} \int \frac{d^3k_1 d\omega_1}{(2\pi)^3} \frac{E_{k_1j}}{\omega_1 + i\delta} \delta(\omega - \omega_1 - \vec{k} \cdot \vec{v}_a + \vec{k}_1 \cdot \vec{v}_a) \\
& \times \left\{ \delta_{ij} + \frac{k_j v_{ai} + k_{1i}v_{aj}}{\omega - \vec{k} \cdot \vec{v}_a + i\delta} + \frac{\vec{k} \cdot \vec{k}_1 v_{ai}v_{aj}}{(\omega - \vec{k} \cdot \vec{v}_a + i\delta)^2} - \frac{v_{ai}v_{aj}}{c^2} \right. \\
& \left. - \frac{\vec{k}_1 \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta)c^2} - \frac{\omega_1}{c^2} \frac{\vec{k} \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a + i\delta)^2} \right\}.
\end{aligned} \quad (70)$$

Changing the integration variable  $k_1 = (\omega_1, \vec{k}_1)$  to  $-k_1$ , equation (70) becomes

$$\begin{aligned}
j_{aki}^{(2)} = & \frac{ie_a^2}{\gamma_a^m} \int \frac{d^3k_1 d\omega_1}{(2\pi)^3} \frac{E_{-k_1j}}{-\omega_1 + i\delta} \delta(\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_a) \\
& \times \left\{ \delta_{ij} + \frac{k_j v_{ai} - k_{1i}v_{aj}}{\omega - \vec{k} \cdot \vec{v}_a + i\delta} - \frac{\vec{k} \cdot \vec{k}_1 v_{ai}v_{aj}}{(\omega - \vec{k} \cdot \vec{v}_a + i\delta)^2} - \frac{v_{ai}v_{aj}}{c^2} \right. \\
& \left. - \frac{\vec{k}_1 \cdot \vec{v}_a v_{ai}v_{aj}}{c^2(\omega_1 - \vec{k}_1 \cdot \vec{v}_a - i\delta)} + \frac{\omega_1}{c^2} \frac{\vec{k} \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a - i\delta)^2} \right\}.
\end{aligned} \quad (71)$$

Combining terms and using the property of the delta function, one notes that

$$\begin{aligned}
& \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_a - \vec{k}_1 \cdot \vec{v}_a) \left[ -\frac{v_{ai}v_{aj}}{c^2} - \frac{\vec{k}_1 \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a - i\delta)c^2} \right. \\
& \left. + \frac{\omega_1}{c^2} \frac{\vec{k} \cdot \vec{v}_a v_{ai}v_{aj}}{(\omega_1 - \vec{k}_1 \cdot \vec{v}_a - i\delta)^2} \right] = \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_a - \vec{k}_1 \cdot \vec{v}_a) \frac{v_{ai}v_{aj} \omega \omega_1}{c^2(\omega - \vec{k} \cdot \vec{v}_a + i\delta)^2}.
\end{aligned} \quad (72)$$

Substituting equation (72) in equation (71) and using the reality property of the field ( $\vec{E}_k = \vec{E}_{-k}^*$ ), then

$$j_{\alpha ki}^{(2)} = -ie_{\alpha} \int \frac{dk_1}{(2\pi)^3} \frac{E_{k_1 j}^*}{\omega_1 - i\delta} \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_{\alpha} - \vec{k}_1 \cdot \vec{v}_{\alpha}) \Lambda_{ij}^{(\alpha)*}(k_1, k) , \quad (73)$$

where

$$\Lambda_{ij}^{(\alpha)}(k_1, k) = \frac{e_{\alpha}}{\gamma_{\alpha} m_{\alpha}} \left\{ \delta_{ij} + \frac{v_{\alpha i} k_j - k_{1i} v_{\alpha j}}{\omega - \vec{k} \cdot \vec{v}_{\alpha} - i\delta} - \frac{v_{\alpha i} v_{\alpha j} (\vec{k} \cdot \vec{k}_1 - \frac{\omega \omega_1}{c^2})}{(\omega - \vec{k} \cdot \vec{v}_{\alpha} - i\delta)^2} \right\} . \quad (74)$$

The currents  $j_{\alpha ki}^{(1)}$  and  $j_{\alpha ki}^{(2)}$  given by equations (64) and (73) are the currents associated with the unperturbed motion and the perturbation in the motion, respectively, of the bare particle. By the same arguments leading to equation (21), the components of the field due to the unperturbed and perturbed test particle trajectory are given by

$$E_{kn}^{(1)} = - \frac{i}{\omega + i\delta} G_{nm} j_{km}^{(1)} \quad (75)$$

and

$$E_{kn}^{(2)} = - \frac{i}{\omega + i\delta} G_{nm} j_{km}^{(2)} , \quad (76)$$

respectively. Therefore, substituting equations (64) and (73) in equations (75) and (76), one obtains

$$E_{kn}^{(1)} = - \frac{ie_{\alpha}}{(2\pi)^3(\omega + i\delta)} v_{\alpha m} G_{nm}(k) \delta(\omega - \vec{k} \cdot \vec{v}_{\alpha}) \quad (77)$$

and

$$E_{kn}^{(2)} = - \frac{e_{\alpha}}{(2\pi)^3(\omega + i\delta)} G_{nm}(k) \int \frac{dk_1}{\omega_1 - i\delta} \frac{E_{k_1 j}^*}{\omega_1 - i\delta} \Lambda_{mj}^{(\alpha)*}(k_1, k) \delta(\omega + \omega_1 - \vec{k} \cdot \vec{v}_{\alpha} - \vec{k}_1 \cdot \vec{v}_{\alpha}) . \quad (78)$$

Equations (77) and (78) agree with equations (16) and (17) of Akopyan and Tsytovich, 1975,<sup>1</sup> except for erroneous overall factors of  $(2\pi)^{-3}$  and  $(2\pi)^{-3}i$  appearing there in equations (16) and (17), respectively. The indices  $n$  and  $m$  are interchanged there as they are in the defining relation, equation (14) of Akopyan,<sup>1</sup> compared to equation (21) here. However, a spatially isotropic plasma is assumed there, so according to equation (22) the Green's function is symmetric, namely  $G_{mn} = G_{nm}$ . Also in Akopyan,<sup>1</sup> single-wave particle resonance is ignored, in which case the small imaginary part  $i\delta$  in equation (74) is

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

ignorable,  $\Lambda_{ij}^{(a)}$  becomes effectively real, and the complex conjugate sign in equation (78) may be removed. The additional factors of  $(4\pi)(2\pi)^3$  in equation (16) and  $(4\pi)$  in (17) there are due to the Gaussian form of Maxwell's equation used there compared to the MKS form used here, and to the difference in Fourier transform convention. For example, in equation (5) of Akopyan and Tsytovich, 1975,<sup>1</sup> the Fourier transform convention employed has a factor of  $(2\pi)^{-3}$  in the inverse Fourier transform in the integration over the three-dimensional wave vector space, and a factor of 1 for the integration over frequency, giving a total factor of  $(2\pi)^{-3}$ . In this report, a total factor of 1 is used, as in equation (7), for example. Likewise, the Fourier transform itself then has a factor of  $(2\pi)^{-1}$  there and  $(2\pi)^{-4}$  here, as in equation (61), for example.

## 5. THE FIELD DUE TO THE DYNAMIC POLARIZATION CURRENT

The Fourier transform of the dynamic polarization current to third-order induced in the beam-plasma is given by<sup>1,\*</sup>

$$\vec{j}_{dpk} = \vec{j}_{dpk}^{(1)} + \vec{j}_{dpk}^{(2)} + \vec{j}_{dpk}^{(3)} \quad (79)$$

The linear polarization current density  $\vec{j}_{dpk}^{(1)}$  is given by

$$j_{dpki}^{(1)} = \sum_s \sigma_{il}^{(s)}(k) E_{kl} \quad (80)$$

Its effects are manifest in the linear Green's function for the beam-plasma system as discussed in section 2. The second-order and third-order polarization current densities  $j_{dpk}^{(2)}$  and  $j_{dpk}^{(3)}$  are given by

$$j_{dpki}^{(2)} = - \sum_s e_s \int \frac{dk_1 dk_2 \delta(k - k_1 - k_2)}{(\omega_1 + i\delta)(\omega_2 + i\delta)} S_{ijl}^{(s)}(k, k_1, k_2) E_{k_1j} E_{k_2l} \quad (81)$$

and

$$j_{dpki}^{(3)} = - \sum_s e_s \int \frac{dk_1 dk_2 dk_3 \delta(k - k_1 - k_2 - k_3)}{(\omega_1 + i\delta)(\omega_2 + i\delta)(\omega_3 + i\delta)} \quad (82)$$

$$\times \Sigma_{ijlm}^{(s)}(k, k_1, k_2, k_3) E_{k_1j} E_{k_2l} E_{k_3m} \quad ,$$

respectively. The second-order nonlinear and third-order nonlinear conductivities for species  $s$ ,  $S_{ijl}^{(s)}$ , and  $\Sigma_{ijlm}^{(s)}$ , respectively, are given by

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

\*H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

$$S_{ijl}^{(s)}(k, k_1, k_2) = e_s^2 \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} \left[ (\omega_1 - \vec{k}_1 \cdot \vec{v}_s) \frac{\partial}{\partial p_{sj}} + v_{jk_1m} \frac{\partial}{\partial p_{sm}} \right] \left[ \frac{\partial}{\partial p_{sl}} + \frac{v_{sl}}{\omega_2 - \vec{k}_2 \cdot \vec{v}_s + i\delta} k_{2n} \frac{\partial}{\partial p_{sn}} \right] f_{ps}^{R(0)}, \quad (83)$$

and

$$\begin{aligned} \Sigma_{ijlm}^{(s)}(k, k_1, k_2, k_3) = & -ie_s^3 \int \frac{d^3 \vec{p}_s}{(2\pi)^3} \frac{v_{si}}{\omega - \vec{k} \cdot \vec{v}_s + i\delta} [\delta_{jn}(\omega_1 - \vec{k}_1 \cdot \vec{v}_s) + k_{1n} v_{sj}] \\ & \times \frac{\partial}{\partial p_{sn}} \frac{1}{\omega - \omega_1 - (\vec{k} - \vec{k}_1) \cdot \vec{v}_s + i\delta} [\delta_{lu}(\omega_2 - \vec{k}_2 \cdot \vec{v}_s) \\ & + k_{2u} v_{sl}] \frac{\partial}{\partial p_{su}} \frac{1}{\omega_3 - \vec{k}_3 \cdot \vec{v}_s + i\delta} [\delta_{mq}(\omega_3 - \vec{k}_3 \cdot \vec{v}_s) \\ & + k_{3q} v_{sm}] \frac{\partial}{\partial p_{sq}} f_{ps}^{R(0)}. \end{aligned} \quad (84)$$

The function  $f_{ps}^{R(0)}$  is the assumed slowly varying and spatially independent background distribution function for species  $s$ , whose Fourier transform is

$$f_k^R(s) = f_{ps}^{R(0)} \delta(k). \quad (85)$$

To determine the fields defined in equation (2) and associated with the respective parts of the dynamic polarization current, one proceeds as in obtaining equations (75) and (76). Thus, one has for the fields produced by the second- and third-order nonlinear dynamic polarization currents, respectively,

$$E_{dpkn}^{(1)} = - \frac{i}{\omega + i\delta} G_{nm}(k) j_{dpkm}^{(2)} \quad (86)$$

and

$$E_{dpkn}^{(2)} = - \frac{i}{\omega + i\delta} G_{nm}(k) j_{dpkm}^{(3)}. \quad (87)$$

Next, substituting equation (81) in equation (86), one obtains

$$\begin{aligned} E_{dpkn}^{(1)} = & \frac{i}{\omega + i\delta} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2 \delta(k - k_1 - k_2)}{(\omega_1 + i\delta)(\omega_2 + i\delta)} \\ & \times S_{mj\ell}^{(s)}(k, k_1, k_2) E_{k_1j} E_{k_2\ell}. \end{aligned} \quad (88)$$

First symmetrizing equation (88) in  $(j, k_1)$  and  $(l, k_2)$ , changing variables of integration from  $k_1$  to  $-k_1$  and  $k_2$  to  $-k_2$ , and using the reality property of the field, then equation (88) becomes

$$E_{dpkn}^{(1)} = \frac{i}{2(\omega + i\delta)} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2}{(\omega_1 - i\delta)(\omega_2 - i\delta)} \delta(k + k_1 + k_2) \quad (89)$$

$$\times \left[ S_{mj\ell}^{(s)}(k, -k_1, -k_2) + S_{m\ell j}^{(s)}(k, -k_2, -k_1) \right] E_{k_1 j}^* E_{k_2 \ell}^* .$$

Comparing equation (89) with equation (22) of Akopyan,<sup>1</sup> a disparity of a factor of  $(1/2)(4\pi)(2\pi)^{-6}$  in the latter is apparently due to explicit inclusion there of a self-field factor of  $(1/2)$ , the use of Gaussian units, and the different Fourier transform and normalization conventions chosen there. The factor of  $(1/2)$  is either a typographical error or results from explicit inclusion of a factor of  $1/2$  associated with self-fields. For the generic expression, equation (89), it is preferable that such a factor be maintained implicit until the formula is applied explicitly and a self-field factor of  $(1/2)$  is needed. The factor of  $(4\pi)$  is due to the use of Gaussian units there. One of the factors of  $(2\pi)^{-3}$  is due to the differing Fourier transform convention. Because of the different Fourier transform convention, the counterpart of equation (81) would have an additional factor of  $(2\pi)^{-3}$ . The other factor of  $(2\pi)^{-3}$  is apparently due to the different normalization of  $f_{ps}^{R(0)}$  there. In short, the  $f_{ps}^{R(0)}$  there is evidently  $(2\pi)^3$  times that here. Alternatively, if the normalization is the same as that here then there is an erroneous factor of  $(2\pi)^{-3}$  appearing there. An isotropic plasma is assumed there, in which case by equation (22) the Green's function is symmetric, namely,  $G_{mn} = G_{nm}$ . However as already noted the indices of  $G_{ij}$  are interchanged in Akopyan and Tsytovich<sup>1</sup> in the definition of the Green's function. It appears, however, that there are typographical errors there. The complex conjugate sign on the fields and a factor of  $e_s$  are left out. Also, there in the second-order nonlinear conductivity, one notes from equation (83) that\*

$$S_{mj\ell}^{(s)}(k, -k_1, -k_2) = S_{mj\ell}^{(s)*}(-k, k_1, k_2) = -\bar{S}_{mj\ell}^{(s)*}(k, k_1, k_2) , \quad (90)$$

where  $\bar{S}_{mj\ell}^{(s)}(k, k_1, k_2)$  designates the conductivity tensor appearing in Akopyan and Tsytovich.<sup>1,\*</sup> In the last step of equation (90) it is recognized that  $S_{mj\ell}^{(s)}(k, k_1, k_2)$  differs from  $S_{mj\ell}^{(s)}(k, k_1, k_2)$  here in that the first complex denominator  $\omega - \vec{k} \cdot \vec{v}_s + i\delta$  in equation (83) here is implicitly  $\omega - \vec{k} \cdot \vec{v}_s - i\delta$

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

\*H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

there.<sup>8,9,\*</sup> Thus, there is also a disagreement in overall sign between equation (89) here and equation (22) of Akopyan and Tsytovich.<sup>1</sup> In summary then, equation (22) of Akopyan and Tsytovich<sup>1</sup> is in error; it should have an additional factor of  $(-e_s)$ , the fields and the conductivity tensor on the right-hand side should be complex conjugated, and the explicit additional factor of  $(1/2)$  there should be omitted.

Next, substituting equation (82) in equation (87),

$$E_{dpkn}^{(2)} = \frac{i}{\omega + i\delta} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3 \delta(k - k_1 - k_2 - k_3)}{(\omega_1 + i\delta)(\omega_2 + i\delta)(\omega_3 + i\delta)} \times \sum_{mijl}^{(s)}(k, k_1, k_2, k_3) E_{k,i} E_{k_2,j} E_{k_3,l} \quad (91)$$

Symmetrizing equation (91) in  $(i, k_1)$ ,  $(j, k_2)$ ,  $(l, k_3)$ , changing variables of integration from  $k_1$  to  $-k_1$ ,  $k_2$  to  $-k_2$ , and  $k_3$  to  $-k_3$ , and using the reality property of the field ( $\vec{E}_{-k} = \vec{E}_k^*$ ), then equation (91) becomes

$$E_{dpkn}^{(2)} = \frac{-i}{6(\omega + i\delta)} G_{nm}(k) \sum_s e_s \int \frac{dk_1 dk_2 dk_3 \delta(k + k_1 + k_2 + k_3)}{(\omega_1 - i\delta)(\omega_2 - i\delta)(\omega_3 - i\delta)} \times \left[ \sum_{mijl}^{(s)}(k, -k_1, -k_2, -k_3) + \sum_{milj}^{(s)}(k, -k_1, -k_3, -k_2) + \sum_{mjil}^{(s)}(k, -k_2, -k_1, -k_3) + \sum_{mjli}^{(s)}(k, -k_2, -k_3, -k_1) + \sum_{mlij}^{(s)}(k, -k_3, -k_1, -k_2) + \sum_{mlji}^{(s)}(k, -k_3, -k_2, -k_1) \right] E_{k,i}^* E_{k_2,j}^* E_{k_3,l}^* \quad (92)$$

Comparing equation (92) with equation (23) of Akopyan and Tsytovich,<sup>1</sup> the disparity of a factor of  $(1/2)(4\pi)(2\pi)^{-9}$  in the latter is again due to an explicit self-field factor of  $(1/2)$ , the use of Gaussian units, and the different Fourier transform and background normalization conventions there.

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

<sup>8</sup>H. E. Brandt, *On the Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma*, Harry Diamond Laboratories, HDL-TR-1970 (February 1982).

<sup>9</sup>H. E. Brandt, *Comment on "Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma,"* *Phys. Fluids* 25 (1982), 1922.

\*H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

Alternatively if the normalization is instead the same as that here, then there is an erroneous factor of  $(2\pi)^{-3}$  appearing there. There is apparently an error there. A factor of  $ie_s$  has been left out. Note that by equation (84) it follows that

$$\Sigma_{mijl}^{(s)}(k, -k_1, -k_2, -k_3) = -\Sigma_{mijl}^{(s)*}(-k, k_1, k_2, k_3) . \quad (93)$$

Also in the conductivity tensor  $T_{ijlm}^{(s)}(k, k_1, k_2, k_3)$  given by equation (21) of Akopyan and Tsytovich,<sup>1</sup> the first two complex denominators are implicitly  $\omega - \vec{k} \cdot \vec{v}_s - i\delta$  and  $\omega + \omega_1 - (\vec{k} + \vec{k}_1) \cdot \vec{v}_s - i\delta$ , respectively, there.\* Also, there is no overall factor of  $i$  as there is in equation (84) here. Therefore,

$$\Sigma_{mijl}^{(s)}(-k, k_1, k_2, k_3) = -iT_{mijl}^{(s)}(k, k_1, k_2, k_3) . \quad (94)$$

Combining equations (93) and (94) then

$$\Sigma_{mijl}^{(s)}(k, -k_1, -k_2, -k_3) = -iT_{mijl}^{(s)*}(k, k_1, k_2, k_3) . \quad (95)$$

In summary then, equation (23) of Akopyan and Tsytovich<sup>1</sup> is also in error and should also have an additional factor of  $(ie_s)$ , the complex conjugate of the fourth-order conductivity tensor there must be taken, and the explicit additional factor of  $(1/2)$  there should be omitted.

## 6. THE TOTAL FIELD

Collecting equations (1), (2), and (3) the total field involved in the bremsstrahlung scattering process and acting on the test particle is given by

$$\vec{E}_k = \vec{E}_k^{(0)} + \vec{E}_k^{(1)} + \vec{E}_k^{(2)} + \vec{E}_{dpk}^{(1)} + \vec{E}_{dpk}^{(2)} . \quad (96)$$

Here  $\vec{E}_k^{(0)}$  is the bremsstrahlung field whose dispersion relations are given by equations (4) and (29) in the spatially isotropic case and equations (4), (20), (38), and (39) in the case of isotropy in both ordinary space and momentum space. The stochastic properties are given by equations (49) and (50). The fields  $\vec{E}_k^{(1)}$  and  $\vec{E}_k^{(2)}$  given by equations (77) and (78) are the self-fields arising from the relativistic test particle's own motion. The fields  $\vec{E}_{dpk}^{(1)}$  and  $\vec{E}_{dpk}^{(2)}$  given by equations (89) and (91) are the increasing order fields produced by the nonlinear dynamic polarization current induced by the test particle. The fields  $\vec{E}_{dpk}^{(1)}$  and  $\vec{E}_{dpk}^{(2)}$  are excited in nonlinear scattering of the field by the induced dynamic polarization current. They involve the three- and four-plasmon vertex, respectively, and play a key role in determining the nonlinear contribution to the collective bremsstrahlung process.

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

\*H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, *Harry Diamond Laboratories*, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

## 7. CONCLUSION

An expression has been obtained, equations (96), (77), (78), (89), and (91), for the total field involved in the collective scattering and bremsstrahlung of a relativistic test particle in a nonequilibrium beam-plasma system. Dispersion relations, equations (4), (29), (20), (38), and (39), and stochastic properties, equations (49) and (50), for the bremsstrahlung field have also been obtained. These results are in agreement with those of Akopyan and Tsytovich<sup>1</sup> with the exception of apparent typographical errors, and have been used by them in their work on nonlinear bremsstrahlung in nonequilibrium plasmas.

The present work, together with related work by the author,<sup>10-14,\*†‡§</sup> is important for ongoing work in calculating collective radiation processes and conditions for the occurrence of radiative instability in relativistic nonequilibrium beam-plasma systems.

---

<sup>1</sup>A. V. Akopyan and V. N. Tsytovich, *Bremsstrahlung in a Nonequilibrium Plasma*, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].

<sup>10</sup>H. E. Brandt, *Symmetries of the Nonlinear Conductivity for a Relativistic Turbulent Plasma*, Harry Diamond Laboratories, HDL-TR-1927 (March 1981).

<sup>11</sup>H. E. Brandt, *Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma*, *Phys. Fluids*, 24 (1981), 1760.

<sup>12</sup>H. E. Brandt, *Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma*, in *Plasma Astrophysics, Course and Workshop Organized by the International School of Plasma Physics*, 27 August to 7 September 1981, Varenna (Como), Italy (European Space Agency ESA SP-161, November 1981), 361 (also to be published by Pergamon Press).

<sup>13</sup>H. E. Brandt, *Symmetry of the Complete Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma*, *Journal of Math. Phys.* 24, (1983), 1332, 2250.

<sup>14</sup>H. E. Brandt, *The Gluckstern-Hull Formula for Electron-Nucleus Bremsstrahlung*, Harry Diamond Laboratories, HDL-TR-1884 (May 1980).

\*H. E. Brandt, *Nonlinear Force on an Unpolarized Relativistic Test Particle to Second Order in the Total Field in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-7 (May 1982) (to be published as HDL-TR-1995, 1983).

†H. E. Brandt, *Nonlinear Dynamic Polarization Force on a Relativistic Test Particle in a Nonequilibrium Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-82-6 (May 1982) (to be published as HDL-TR-1994, 1983).

‡H. E. Brandt, *Theoretical Methods in the Calculation of the Bremsstrahlung Recoil Force in a Nonequilibrium Relativistic Beam-Plasma System*, Harry Diamond Laboratories, HDL-PRL-83-6 (April 1983) (to be published as HDL-TR-2009, 1983).

§Other related work prepared in preprint form will be published later and is available from the author.

# LITERATURE CITED

- (1) A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung in a Nonequilibrium Plasma, *Fiz. Plazmy*, 1 (1975), 673 [*Sov. J. Plasma Phys.*, 1 (1975), 371].
- (2) A. V. Akopyan and V. N. Tsytovich, Transition Bremsstrahlung of Relativistic Particles, *Zh. Eksp. Teor. Fiz.*, 71 (1976), 166 [*Sov. Phys. JETP*, 44 (1976), 87].
- (3) A. V. Akopyan and V. N. Tsytovich, Bremsstrahlung Instability of Relativistic Electrons in a Plasma, *Astrofizika*, 13 (1977), 717 [*Astrophysics*, 13 (1977), 423].
- (4) V. N. Tsytovich, Collective Effects in Bremsstrahlung of Fast Particles in Plasmas, *Comments Plasma Phys. Conf., Fusion*, 4 (1978), 73.
- (5) A. I. Akhiezer, I. A. Akhiezer, R. V. Polovin, A. G. Sitenko, and K. N. Stepanov, *Plasma Electrodynamics*, 1, Linear Theory, Pergamon Press (1975), 206.
- (6) V. N. Tsytovich, *Nonlinear Effects in Plasma*, Plenum Press, New York (1970), 31.
- (7) V. N. Tsytovich, *Theory of Turbulent Plasma* (Consultants Bureau), Plenum Press, New York (1977), 63-65.
- (8) H. E. Brandt, On the Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, Harry Diamond Laboratories, HDL-TR-1970 (February 1982).
- (9) H. E. Brandt, Comment on "Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma," *Phys. Fluids* 25 (1982), 1922.
- (10) H. E. Brandt, Symmetries of the Nonlinear Conductivity for a Relativistic Turbulent Plasma, Harry Diamond Laboratories, HDL-TR-1927 (March 1981).
- (11) H. E. Brandt, Exact Symmetry of the Second-Order Nonlinear Conductivity for a Relativistic Turbulent Plasma, *Phys. Fluids*, 24 (1981), 1760.
- (12) H. E. Brandt, Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, in *Plasma Astrophysics, Course and Workshop Organized by the International School of Plasma Physics*, 27 August to 7 September 1981, Varenna (Como), Italy (European Space Agency ESA SP-161, November 1981) 361 (also to be published by Pergamon Press).

LITERATURE CITED (Cont'd)

- (13) H. E. Brandt, Symmetry of the Complete Second-Order Nonlinear Conductivity Tensor for an Unmagnetized Relativistic Turbulent Plasma, J. Math. Phys. 24, (1983), 1332, 2250.
- (14) H. E. Brandt, The Gluckstern-Hull Formula for Electron-Nucleus Bremsstrahlung, Harry Diamond Laboratories, HDL-TR-1884 (May 1980).

## DISTRIBUTION

ADMINISTRATOR  
DEFENSE TECHNICAL INFORMATION CENTER  
ATTN DTIC-DDA (12 COPIES)  
CAMERON STATION, BUILDING 5  
ALEXANDRIA, VA 22314

COMMANDER  
US ARMY MATERIEL DEVELOPMENT & READINESS  
COMMAND  
5001 EISENHOWER AVENUE  
ALEXANDRIA, VA 22333

DIRECTOR  
DEFENSE ADVANCED RESEARCH PROJECTS  
AGENCY  
ATTN R. GULLICKSON  
ATTN L. C. MARQUET  
ATTN R. SEPUCHA  
1400 WILSON BLVD  
ARLINGTON, VA 22209

DIRECTOR  
DEFENSE COMMUNICATIONS AGENCY  
WASHINGTON, DC 20305

DIRECTOR  
DEFENSE INTELLIGENCE AGENCY  
ATTN D. SPOHN  
ATTN DT-1, NUCLEAR & APPLIED  
SCIENCES DIV  
ATTN ELECTRONIC WARFARE BRANCH  
WASHINGTON, DC 20301

UNDER SECRETARY OF DEFENSE  
FOR RESEARCH & ENGINEERING  
ATTN DEPUTY UNDER SECRETARY  
(RESEARCH & ADVANCED TECH)  
ATTN D. L. LAMBERSON  
WASHINGTON, DC 20301

OFFICE OF THE UNDER SECRETARY OF  
DEFENSE FOR RESEARCH & ENGINEERING  
DIR ENERGY TECHNOLOGY OFFICE  
THE PENTAGON  
WASHINGTON, DC 20301

DIRECTOR  
DEFENSE NUCLEAR AGENCY  
ATTN DEPUTY DIRECTOR  
SCIENTIFIC TECHNOLOGY  
ATTN G. BAKER  
ATTN RAEV, ELECTRONIC VULNERABILITY  
ATTN J. Z. FARBER  
ATTN G. K. SOPER  
ATTN V. VAN LINT  
WASHINGTON, DC 20305

DIRECTOR  
NATIONAL SECURITY AGENCY  
ATTN TECHNICAL LIBRARY  
ATTN F. BEDARD  
FT MEADE, MD 20755

ASSISTANT SECRETARY OF THE ARMY (RDA)  
ATTN DEP FOR SCI & TECH  
WASHINGTON, DC 20301

OFFICE, DEPUTY CHIEF OF STAFF  
FOR OPERATIONS & PLANS  
DEPT OF THE ARMY  
ATTN DAMO-SSN, NUCLEAR DIV  
WASHINGTON, DC 20310

OFFICE OF THE DEPUTY CHIEF OF STAFF  
FOR RESEARCH, DEVELOPMENT,  
& ACQUISITION  
DEPARTMENT OF THE ARMY  
ATTN DIRECTOR OF ARMY RESEARCH,  
M. E. LASSER  
ATTN DAMA-CSS-N, NUCLEAR TEAM  
ATTN DAMA-ARZ-O, F. D. VERDERAME  
WASHINGTON, DC 20310

COMMANDER  
US ARMY ABERDEEN PROVING GROUND  
ATTN STEAP-TL, TECH LIB  
ABERDEEN PROVING GROUND, MD 21005

COMMANDER  
BALLISTIC MISSILE DEFENSE  
ADVANCED TECHNOLOGY CENTER  
ATTN D. SCHENK  
PO BOX 1500  
HUNTSVILLE, AL 35800

DIRECTOR  
US ARMY BALLISTIC RESEARCH LABORATORY  
ATTN DRDAR-TSB-S (STINFO)  
ATTN D. ECCLESHALL  
ATTN C. HOLLANDSWORTH  
ATTN R. KREMENS  
ABERDEEN PROVING GROUND, MD 21005

US ARMY ELECTRONICS TECHNOLOGY &  
DEVICES LABORATORY  
ATTN DELET-DD  
ATTN N. WILSON  
FT MONMOUTH, NJ 07703

US ARMY ERADCOM  
ATTN C. M. DESANTIS  
FT MONMOUTH, NJ 07703

DISTRIBUTION (Cont'd)

COMMANDER  
US ARMY FOREIGN SCIENCE  
& TECHNOLOGY CENTER  
FEDERAL OFFICE BLDG  
ATTN DRXST-SD, SCIENCES DIV  
ATTN T. CALDWELL  
220 7TH STREET, NE  
CHARLOTTESVILLE, VA 22901

DIRECTOR  
US ARMY MATERIEL SYSTEMS  
ANALYSIS ACTIVITY  
ATTN DRXSY-MP  
ABERDEEN PROVING GROUND, MD 21005

COMMANDER  
US ARMY MATERIALS & MECHANICS  
RESEARCH CENTER  
ATTN DRXMR-H, BALLISTIC MISSILE  
DEF MATLS PROG OFC  
WATERTOWN, MA 02172

COMMANDER  
US ARMY MISSILE COMMAND  
ATTN DRDMI-TR, PHYSICAL SCIENCES DIR  
REDSTONE ARSENAL, AL 35809

COMMANDER  
US ARMY MISSILE & MUNITIONS  
CENTER & SCHOOL  
ATTN ATSK-CTD-F  
REDSTONE ARSENAL, AL 35809

ARMY RESEARCH OFFICE (DURHAM)  
PO BOX 12211  
ATTN H. ROBL  
ATTN R. LONTZ  
ATTN B. D. GUENTHER  
ATTN TECH LIBRARY  
RESEARCH TRIANGLE PARK, NC 27709

COMMANDER  
US ARMY RSCH & STD GP (EUR)  
ATTN CHIEF, PHYSICS & MATH BRANCH  
FPO NEW YORK 09510

NAVAL AIR SYSTEMS COMMAND  
ATTN LCDR G. BATES, PMS-405  
WASHINGTON, DC 20361

NAVAL INTELLIGENCE SUPPORT CENTER  
ATTN M. KOONTZ  
4301 SUITLAND RD  
SUITLAND, MD 20390

NAVAL MATERIAL COMMAND  
ATTN T. HORWATH  
2211 JEFFERSON DAVIS HWY  
WASHINGTON, DC 20301

OFFICE OF NAVAL RESEARCH  
ATTN C. ROBERSON  
ATTN W. J. CONDELL  
800 N. QUINCY ST  
ARLINGTON, VA 22217

SUPERINTENDENT  
NAVAL POSTGRADUATE SCHOOL  
ATTN LIBRARY, CODE 2124  
MONTEREY, CA 93940

DIRECTOR  
NAVAL RESEARCH LABORATORY  
ATTN 2600, TECHNICAL INFO DIV  
ATTN 5540, LASER PHYSICS  
ATTN 6000, MATL & RADIATION SCI & TE  
ATTN B. RIPIN  
ATTN L. A. COSBY  
ATTN E. E. KEMPE  
ATTN J. T. SCHRIEMPF  
ATTN R. F. WENZEL  
ATTN R. HETTCHE  
ATTN J. GOLDEN  
ATTN V. L. GRANATSTEIN  
ATTN M. A. REED  
ATTN R. K. PARKER  
ATTN P. A. SPRANGLE  
ATTN S. GOLD  
ATTN C. A. KAPETANAKOS  
ATTN S. AMN  
ATTN T. P. COFFEY  
ATTN R. JACKSON  
ATTN I. M. VITKOVITSKY  
ATTN M. FRIEDMAN  
ATTN J. PASOUR  
ATTN G. COOPERSTEIN  
WASHINGTON, DC 20375

COMMANDER  
NAVAL SURFACE WEAPONS CENTER  
ATTN V. PUGLIELLI  
ATTN DX-21, LIBRARY DIV  
DAHLGREN, VA 22448

COMMANDER  
NAVAL SURFACE WEAPONS CENTER  
ATTN J. Y. CHOE  
ATTN F. SAZAMA  
ATTN H. UHM  
ATTN V. KENYON  
ATTN E. NOLTING  
ATTN WA-13, HIGH-ENERGY LASER BR  
ATTN WA-50, NUCLEAR WEAPONS  
EFFECTS DIV  
ATTN WR, RESEARCH & TECHNOLOGY DEPT  
ATTN WR-40, RADIATION DIV  
ATTN E-43, TECHNICAL LIB  
WHITE OAK, MD 20910

DISTRIBUTION (Cont'd)

COMMANDER  
NAVAL WEAPONS CENTER  
ATTN 315, LASER/INFRARED SYS DIV  
ATTN 381, PHYSICS DIV  
CHINA LAKE, CA 93555

HQ USAF/SAMI  
WASHINGTON, DC 20330

ASSISTANT SECRETARY OF THE AIR FORCE  
(RESEARCH & DEVELOPMENT)  
WASHINGTON, DC 20330

DIRECTOR  
AF OFFICE OF SCIENTIFIC RESEARCH  
BOLLING AFB  
ATTN NP, DIR OF PHYSICS  
ATTN M. A. STROSCIO  
ATTN R. BARKER  
WASHINGTON, DC 20332

COMMANDER  
AF WEAPONS LAB, AFSC  
ATTN J. GENEROSA  
ATTN A. H. GUENTHER  
ATTN W. E. PAGE  
ATTN LR, LASER DEV DIV  
KIRTLAND AFB, NM 87117

WRIGHT-PATTERSON AFB  
FOREIGN TECHNOLOGY DIVISION/ETD  
ATTN J. BUTLER  
WRIGHT-PATTERSON AFB, OH 45433

CENTRAL INTELLIGENCE AGENCY  
ATTN R. PETTIS  
ATTN D. B. NEWMAN  
PO BOX 1925  
WASHINGTON, DC 20013

DEPARTMENT OF COMMERCE  
NATIONAL BUREAU OF STANDARDS  
ATTN LIBRARY  
ATTN CENTER FOR RADIATION RESEARCH  
ATTN C. TEAGUE  
ATTN E. MARX  
WASHINGTON, DC 20234

DEPARTMENT OF ENERGY  
ATTN T. F. GODLOVE  
ATTN M. MURPHY  
ATTN A. COLE  
ATTN O. P. MANLEY  
WASHINGTON, DC 20585

NATIONAL AERONAUTICS & SPACE  
ADMINISTRATION  
ATTN R. RAMATY  
GODDARD SPACE FLIGHT CENTER  
GREENBELT, MD 20771

ADMINISTRATOR  
NASA HEADQUARTERS  
WASHINGTON, DC 20546

DIRECTOR  
NASA  
GODDARD SPACE FLIGHT CENTER  
ATTN 250, TECH INFO DIV  
GREENBELT, MD 20771

AMES LABORATORY (ERDA)  
IOWA STATE UNIVERSITY  
ATTN NUCLEAR SCIENCE CATEGORY  
AMES, IA 50011

ASTRONOMICAL INSTITUTE--ZONNENBURG 2  
ATTN M. KUPERUS  
3512 NL UTRECHT, THE NETHERLANDS

THE AUSTRALIAN NATIONAL UNIVERSITY  
DEPT OF THEORETICAL PHYSICS  
ATTN D. B. MELROSE  
PO BOX 4  
CANBERRA A.C.T. 2600  
AUSTRALIA

B. K. DYNAMICS  
ATTN R. J. LINZ  
15825 SHADY GROVE RD  
ROCKVILLE, MD 20850

BOEING AEROSPACE  
ATTN H. T. BUSER  
PO BOX 3999 MS 8R-10  
SEATTLE, WA 98124

BROOKHAVEN NATIONAL LABORATORY  
ASSOCIATED UNIVERSITIES, INC  
ATTN PHYSICS DEPT  
UPTON, LONG ISLAND, NY 11973

CALIFORNIA INSTITUTE OF TECHNOLOGY  
THEORETICAL ASTROPHYSICS  
ATTN K. THORNE  
130-33  
PASADENA, CA 91125

UNIVERSITY OF CALIFORNIA  
DEPT OF APPL PHYS & INFORMATION  
SCIENCE  
ATTN H. ALFVEN  
LA JOLLA, CA 92093

CALIFORNIA INSTITUTE OF TECHNOLOGY  
JET PROPULSION LABORATORY  
ATTN J. C. HUBBARD  
IRVINE, CA 92717

DISTRIBUTION (Cont'd)

UNIVERSITY OF CALIFORNIA  
ELECTRICAL ENGINEERING DEPARTMENT  
ATTN C. K. BIRDSALL  
BERKELEY, CA 94720

UNIVERSITY OF CALIFORNIA  
PHYSICS DEPT  
ATTN A. N. KAUFMAN  
BERKELEY, CA 94720

UNIVERSITY OF CALIFORNIA, DAVIS  
APPLIED SCIENCES  
ATTN J. DEGROOT  
DAVIS, CA 95616

UNIVERSITY OF CALIFORNIA, IRVINE  
DEPT OF PHYSICS  
ATTN G. BENFORD  
ATTN N. ROSTOKER  
ATTN M. MAYER  
ATTN A. RAY  
IRVINE, CA 92717

UNIVERSITY OF CALIFORNIA, LOS ANGELES  
DEPT OF PHYSICS  
ATTN K. NOZAKI  
LOS ANGELES, CA 90025

CAMBRIDGE UNIVERSITY  
INSTITUTE OF ASTRONOMY  
MADINGLEY ROAD  
ATTN M. REES  
CAMBRIDGE CB 3 0 HA, ENGLAND

CHALMERS UNIV OF TECHNOLOGY  
INST OF ELECTROMAGNETIC FIELD  
THEORY  
ATTN H. WILHELMSSON  
S-41296 GOTHENBURG, SWEDEN

UNIVERSITY OF CHICAGO  
LAB FOR ASTROPHYSICS & SPACE RESEARCH  
ATTN E. PARKER  
CHICAGO, IL 60637

UNIVERSITY OF COLORADO  
DEPT OF ASTROGEOPHYSICS  
ATTN M. GOLDMAN  
ATTN D. SMITH  
BOULDER, CO 80309

COLUMBIA UNIVERSITY  
ATTN S. JOHNSTON  
216 MUDD BLDG  
NEW YORK, NY 10027

CORNELL UNIVERSITY  
ATTN R. LOVELACE  
ATTN R. N. SUDAN  
ATTN J. NATION  
ATTN D. HAMMER  
ATTN H. FLEISHMANN  
ITHACA, NY 14853

DARTMOUTH COLLEGE  
PHYSICS DEPT  
ATTN J. E. WALSH  
HANOVER, NH 03755

UNIVERSITY OF THE DISTRICT  
OF COLUMBIA  
VAN NESS CAMPUS  
DEPT OF PHYSICS  
ATTN M. J. SMITH  
4200 CONNECTICUT AVE, NW  
WASHINGTON, DC 20008

ENGINEERING SOCIETIES LIBRARY  
345 EAST 47TH STREET  
ATTN ACQUISITIONS DEPARTMENT  
NEW YORK, NY 10017

GENERAL DYNAMICS  
ATTN K. H. BROWN  
PO BOX 2507 MZ 44-21  
POMONA, CA 91769

UNIVERSITY OF ILLINOIS AT  
URBANA--CHAMPAIGN  
DEPT OF PHYSICS  
ATTN N. IWAMOTO  
URBANA, IL 61801

ISTITUTO DI FISICA DELL' UNIVERSITA  
VIA CELORIA 16  
ATTN P. CALDIROLA  
ATTN C. PAIZIS  
ATTN E. SINDONI  
20133 MILANO, ITALY

ISTITUTO DI FISICA DELL' UNIVERSITA  
ATTN A. CAVALIERE  
ATTN R. RUFFINI  
ROME, ITALY

ISTITUTO DI FISICA GENERALE  
DELL' UNIVERSITA  
CORSO M. D'AZEGLIO  
ATTN A. FERRARI  
46 TORINO, ITALY

JAYCOR  
ATTN E. CONRAD  
205 S. WHITING ST  
ALEXANDRIA, VA 22304

DISTRIBUTION (Cont'd)

KAMAN SCIENCES CORP  
ATTN T. A. TUMOLILLO  
1500 GARDEN OF THE GODS  
COLORADO SPRINGS, CO 80907

LAWRENCE BERKELEY LAB  
ATTN A. FALTENS  
ATTN A. M. SESSLER  
ATTN D. KEEFE  
BERKELEY, CA 94720

LAWRENCE LIVERMORE LABORATORY  
UNIVERSITY OF CALIFORNIA  
ATTN L. MARTIN  
ATTN P. WHEELER  
ATTN H. S. CABAYAN  
ATTN R. BRIGGS  
ATTN E. K. MILLER  
ATTN R. ZIOLKOWSKI  
ATTN R. SCARPETTI  
ATTN R. MINICH  
ATTN S. BURKHART  
ATTN G. VOGTLIN  
ATTN V. W. SLIVINSKY  
ATTN S. L. YU  
ATTN G. CRAIG  
ATTN J. WYATT  
ATTN R. ALVAREZ  
ATTN G. LASCHE  
ATTN J. H. YEE  
ATTN H. W. MELDNER  
ATTN J. S. PETTIBONE  
ATTN A. J. POGGIO  
ATTN M. LOYD  
PO BOX 808  
LIVERMORE, CA 94550

CHIEF  
LIVERMORE DIVISION, FIELD COMMAND, DNA  
LAWRENCE LIVERMORE LABORATORY  
PO BOX 808  
LIVERMORE, CA 94550

LOS ALAMOS SCI LAB  
ATTN S. COLGATE  
ATTN D. F. DUBOIS  
ATTN D. FORSLAND  
ATTN C. M. FOWLER  
ATTN B. WARNER  
ATTN J. LANDT  
ATTN R. HOEBERLING  
ATTN T. R. KING  
ATTN A. KADISH  
ATTN K. LEE  
ATTN R. W. FREYMAN  
ATTN A. W. CHURMATZ  
ATTN L. M. DUNCAN  
ATTN D. B. HENDERSON  
ATTN L. E. THODE  
ATTN H. A. DAVIS  
ATTN T. KWAN  
PO BOX 1663  
LOS ALAMOS, NM 87545

UNIVERSITY OF MARYLAND  
DEPT OF ELECTRICAL ENGINEERING  
ATTN M. REISER  
ATTN W. DESTLER  
ATTN C. D. STRIFFLER  
ATTN M. T. RHEE  
COLLEGE PARK, MD 20742

UNIVERSITY OF MARYLAND  
DEPT OF PHYSICS & ASTRONOMY  
ATTN H. R. GRIEM  
ATTN E. OTT  
ATTN C. GREBOGI  
ATTN K. PAPADOPOULOS  
ATTN J. WEBER  
COLLEGE PARK, MD 20742

MASSACHUSETTS INSTITUTE  
OF TECHNOLOGY  
PLASMA FUSION CENTER  
ATTN G. BEKEFI  
ATTN J. BELCHER  
ATTN T. CHANG  
ATTN B. COPPI  
ATTN R. DAVIDSON  
ATTN C. LIN  
ATTN S. OLBERT  
ATTN B. ROSSI  
CAMBRIDGE, MA 02139

MAX PLANCK INST, FOR AERONOMIE  
3411 KATLENBG  
ATTN J. AXFORD  
ATTN E. MARSCH  
LINDAU 3, WEST GERMANY

MAX PLANCK INSTITUTE OF PHYSICS  
& ASTROPHYSICS  
ATTN M. NORMAN  
ATTN J. TRUMPER  
GARCHING BEI MUNICH, WEST GERMANY

MAXWELL LABORATORIES  
ATTN J. S. PEARLMAN  
8835 BALBOA AVE  
SAN DIEGO, CA 92123

MISSION RESEARCH CORP  
ATTN B. GODFREY  
ATTN D. J. SULLIVAN  
ATTN M. BOLLEN  
ATTN C. LONGMIRE  
ATTN D. VOSS  
1720 RANDOLPH RD, SE  
ALBUQUERQUE, NM 87106

UNIVERSITY OF NAGOYA  
DEPT OF PHYSICS  
ATTN S. HAYAKAWA  
NAGOYA, JAPAN

DISTRIBUTION (Cont'd)

NATIONAL RESEARCH COUNCIL  
DIVISION OF PHYSICS  
ATTN P. JAANIMAGI  
OTTOWA, ONTARIO  
CANADA

NATIONAL SCIENCE FOUNDATION  
DIVISION OF ASTRONOMICAL SCIENCES  
ATTN M. AIZENMAN  
1800 G STREET, NW  
WASHINGTON, DC 20550

OAK RIDGE NATIONAL LABORATORY  
PO BOX Y  
ATTN A. C. ENGLAND  
OAK RIDGE, TN 37830

OCCIDENTAL RESEARCH CORP  
ATTN D. B. CHANG  
2100 SE MAIN  
IRVINE, CA 92713

OXFORD UNIVERSITY  
DEPARTMENT THEORETICAL PHYSICS  
1 KEBLE RD  
ATTN D. TER HAAR  
OXFORD OX1 3NP, ENGLAND

OXFORD UNIVERSITY  
ASTROPHYSICS DEPARTMENT  
SOUTH PARK ROAD  
ATTN A. HALL  
OXFORD--OXI 3 RQ, ENGLAND

OXFORD UNIVERSITY  
DEPT OF ENGINEERING SCIENCE  
PARKS ROAD  
ATTN L. M. WICKENS  
ATTN J. E. ALLEN  
OXFORD, UNITED KINGDOM

PHYSICS INTERNATIONAL  
ATTN B. A. LIPPMANN  
ATTN R. D. GENUARIO  
2700 MERCER ST  
SAN LEANDRO, CA 94577

PRINCETON UNIVERSITY  
ASTROPHYSICAL SCIENCES  
PETON HALL  
ATTN R. KULSRUD  
PRINCETON, NJ 08540

PRINCETON UNIVERSITY  
PLASMAPHYSICS LABORATORY  
ATTN R. WHITE  
PRINCETON, NJ 08540

QUEEN MARY COLLEGE  
DEPT OF APPLIED MATH  
MILE END ROAD  
ATTN D. BURGESS  
LONDON EI 4NS, ENGLAND

RUHR UNIVERSITY  
ATTN W. SEBOLD  
ATTN W. ZWINGMANN  
4630 BOCHUM 1, WEST GERMANY

INSTITUTE OF EXPERIMENTAL PHYSICS  
RUHR-UNIVERSITY  
POSTFACH 2148  
ATTN H. KUNZE  
436 BOCHUM, WEST GERMANY

SACHS/FREEMAN ASSOC  
ATTN R. A. MAHAFFEY  
19300 GALLANT FOX  
SUITE 214  
BOWIE, MD 20715

SANDIA LABORATORIES  
ATTN R. B. MILLER  
ATTN J. POWELL  
ATTN W. BALLARD  
ATTN C. EKDAHL  
ATTN W. D. BROWN  
ATTN S. HUMPHRIES, JR.  
ATTN G. ROHWEIN  
ATTN K. PRESTWICH  
4244 SANDIA LABS  
ALBUQUERQUE, NM 87115

SCIENCE APPLICATIONS, INC  
ATTN E. P. CORNET  
ATTN A. DROBOT  
ATTN E. KANE  
1710 GOODRIDGE DR  
PO BOX 1303  
MCLEAN, VA 22012

SCIENCE APPLICATIONS, INC  
ATTN F. CHILTON  
1200 PROSPECT ST  
PO BOX 2351  
LA JOLLA, CA 92038

SERVICE DE CHEMIE PHYSIQUELL  
ATTN R. BALESCU  
CAMPUS PLAINE U.L.B.  
CODE POSTAL n° 231  
BOULEVARD DU TRIOMPHE  
10 50 BRUXELLES, BELGIUM

SPECOLA VATICANA  
ATTN W. STOEGER, S. J.  
I-00120 CITTA DEL VATICANO  
ITALY

# DISTRIBUTION (Cont'd)

SRI INTERNATIONAL  
ATTN G. AUGUST  
ATTN C. L. RINO  
333 RAVENSWOOD AVE  
MENLO PARK, CA 94025

STANFORD UNIVERSITY  
INST PLASMA RES  
ATTN P. A. STURROCK  
STANFORD, CA 94305

STERRERWACHT-LEIDEN  
ATTN C. A. NORMAN  
ATTN P. ALLAN  
2300 RA LEIDEN, THE NETHERLANDS

TEL-AVIV UNIVERSITY  
DEPT OF PHYSICS & ASTRONOMY  
ATTN G. TAUBER  
TEL AVIV, ISRAEL

TELEDYNE BROWN ENGINEERING  
CUMMINGS RESEARCH PARK  
ATTN MELVIN L. PRICE, MS-44  
HUNTSVILLE, AL 35807

INSTITUTE FOR THEORETICAL MECHANICS-  
RUG  
KRIJGSLAAN 271-S9  
ATTN F. VERHEEST  
B-9000 GENT, BELGIUM

UNIVERSITY OF TENNESSEE  
DEPT OF ELECTRICAL ENGINEERING  
ATTN I. ALEXEFF  
ATTN J. R. ROTH  
KNOXVILLE, TN 37916

UMEA UNIVERSITY  
DEPT OF PLASMA PHYSICS  
ATTN J. LARSSON  
ATTN L. STENFLO  
S-90187 UMEA  
SWEDEN

UNIVERSITY OF WASHINGTON  
DEPT OF PHYSICS  
ATTN M. BAKER  
SEATTLE, WA 98195

WEIZMANN INSTITUT  
DEPT OF NUCLEAR PHYSICS  
ATTN AMRI WANDEL  
REHOVOT, ISRAEL

WESTERN RESEARCH CORP  
ATTN R. O. HUNTER  
225 BROADWAY, SUITE 1600  
SAN DIEGO, CA 92101

US ARMY ELECTRONICS RESEARCH &  
DEVELOPMENT COMMAND  
ATTN COMMANDER, DRDEL-CG  
ATTN TECHNICAL DIRECTOR, DRDEL-CT  
ATTN PUBLIC AFFAIRS OFFICE, DRDEL-IN

HARRY DIAMOND LABORATORIES  
ATTN CO/TD/TSO/DIVISION DIRECTORS  
ATTN RECORD COPY, 81200  
ATTN HDL LIBRARY, 81100 (3 COPIES)  
ATTN HDL LIBRARY (WOODBIDGE), 81100  
ATTN TECHNICAL REPORTS BRANCH, 81300  
ATTN LEGAL OFFICE, 97000  
ATTN CHAIRMAN, EDITORIAL COMMITTEE  
ATTN MORRISON, R. E., 13500  
ATTN CHIEF, 21000  
ATTN CHIEF, 21100  
ATTN CHIEF, 21200  
ATTN CHIEF, 21300  
ATTN CHIEF, 21400  
ATTN CHIEF, 21500  
ATTN CHIEF, 22000  
ATTN CHIEF, 22100  
ATTN CHIEF, 22300  
ATTN CHIEF, 22800  
ATTN CHIEF, 22900  
ATTN CHIEF, 20240  
ATTN CHIEF, 11000  
ATTN CHIEF, 13000  
ATTN CHIEF, 13200  
ATTN CHIEF, 13300  
ATTN CHIEF, 13500  
ATTN CHIEF, 15200  
ATTN BROWN, E., 00211  
ATTN SINDORIS, A., 00211  
ATTN GERLACH, H., 11100  
ATTN LIBELO, L., 11200  
ATTN LOKERSON, D., 11400  
ATTN CROWNE, F., 13200  
ATTN DROPKIN, H., 13200  
ATTN LEAVITT, R., 13200  
ATTN MORRISON, C., 13200  
ATTN SATTLER, J., 13200  
ATTN WORTMAN, D., 13200  
ATTN KULPA, S., 13300  
ATTN SILVERSTEIN, J., 13300  
ATTN FAZI, C., 13500  
ATTN LOMONACO, S., 15200  
ATTN J. CORRIGAN, 20240  
ATTN FARRAR, F., 21100  
ATTN GARVER, R., 21100  
ATTN TATUM, J., 21100  
ATTN MERKEL, G., 21300  
ATTN MCLEAN, B., 22300  
ATTN OLDHAM, T., 22300  
ATTN BLACKBURN, J., 22800  
ATTN GILBERT, R., 22800  
ATTN KLEBERS, J., 22800  
ATTN VANDERWALL, J., 22800  
ATTN BROMBORSKY, A., 22900

DISTRIBUTION (Cont'd)

HARRY DIAMOND LABORATORIES (Cont'd)

ATTN DAVIS, D., 22900  
ATTN GRAYBILL, S., 22900  
ATTN HUTTLIN, G. A., 22900  
ATTN KEHS, A., 22900  
ATTN KERRIS, K., 22900  
ATTN LAMB, R., 22900  
ATTN LINDSAY, D., 22900  
ATTN LITZ, M., 22900  
ATTN RUTH, B., 22900  
ATTN STEWART, A., 22900  
ATTN SOLN, J., 22900  
ATTN WHITTAKER, D., 22900  
ATTN ELBAUM, S., 97100  
ATTN BRANDT, H. E., 22300  
(60 COPIES)

